

Tilburg University

Modification of the Kojima-Nishino-Arima algorithm and its computational complexity

Gao, T.; Talman, A.J.J.; Wang, Z.

Publication date:
1991

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Gao, T., Talman, A. J. J., & Wang, Z. (1991). *Modification of the Kojima-Nishino-Arima algorithm and its computational complexity*. (CentER Discussion Paper; Vol. 1991-27). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

No. 9127

**MODIFICATION OF THE KOJIMA-NISHINO-ARIMA
ALGORITHM AND ITS COMPUTATIONAL COMPLEXITY**

by T. Gao, A.J.J. Talman
and Z. Wang

July 1991

ISSN 0924-7815

MODIFICATION OF THE KOJIMA-NISHINO-ARIMA ALGORITHM AND ITS COMPUTATIONAL COMPLEXITY*

T. Gao[‡] A.J.J. Talman[‡] Z. Wang[†]

May, 1991

Abstract

A modification of the Kojima-Nishino-Arima algorithm for finding all zeros of a polynomial is presented. This modification yields a polynomial-time result on its computational complexity. The main theorem asserts that the cost of locating all zeros of a polynomial of degree n up to an accuracy of ε grows no faster than $O(\max\{n^4, n^3 \log_2(n/\varepsilon)\})$.

Keywords: zeros of polynomials, piecewise linear homotopy, error estimate, computational complexity

*This work was completed when the third author was visiting the Center for Economic Research (CentER), Tilburg University, Tilburg, The Netherlands. This author is grateful to CentER for the invitation and hospitality.

[†]Lingnan College, Zhongshan University, Guangzhou, China

[‡]Department of Econometrics, Tilburg University, Tilburg, The Netherlands

1 Introduction

In response to a question by Smale [6] about the efficiency of piecewise linear homotopy methods in solving polynomial equations, there have been several results, see [3],[5],[9]. However, these results are all dealing with Kuhn's algorithm for polynomials, see [2]. There are two flaws to be indicated. First, as a zero finding method, Kuhn's algorithm fails to give correct multiplicities ("A counterexample for a theorem of Kuhn", presented by Kuhn at the 17th TIMS/ORSA joint National Meeting, San Francisco, 1984). Second, employing integer labelling, Kuhn's algorithm is not, in the original sense of piecewise linear path-following, an exact PL homotopy method.

On the other hand, Kojima, Nishino and Arima [1] presented a PL homotopy algorithm (KNA algorithm in short) employing vector labelling. Due to the introduction of a perturbation term, it can give all zeros of a polynomial with correct multiplicities. This feature makes the algorithm much more attractive. However, there is not any result on its computational complexity yet. It is well-known that, for a complexity discussion of certain kind, an error estimate of the computation is needed first. In spite of the original asymptotic estimate in [1], the desired estimate should be independent from the coefficients of the polynomial. In this paper we obtain an error estimate which is explicit only in terms of the homotopy parameter and of the degree of the polynomial. In doing so, the perturbation term, which makes the KNA algorithm successful, appears to be the problem. The key step of our work is to modify the perturbation. This modification yields a desired estimate. After that, the main result, that the cost of locating all zeros of a polynomial of degree n up to an accuracy of $\varepsilon > 0$ with correct multiplicities grows no faster than $O(\max\{n^4, n^3 \log_2(n/\varepsilon)\})$, follows.

2 Modification of perturbation

Let $f(z) = z^n + \sum_{k=1}^n a_k z^{n-k}$ be a polynomial with complex coefficients. To compute the n roots of $f(z)$ we choose an auxiliary polynomial $h(z) = z^n - r_0^n$, where r_0 is a positive

constant. Let the function $H : [0, 1] \times \mathbb{C} \rightarrow \mathbb{C}$ be defined by

$$H(t, z) = \begin{cases} f(z) + (2t)^{2nb} & \text{if } t \in [0, \frac{1}{2}], \\ (2t - 1)h(z) + 2(1 - t)f(z) + b & \text{if } t \in [\frac{1}{2}, 1], \end{cases} \quad (2.1)$$

where b is a nonzero complex constant. Then H is a homotopy between the polynomial f on level 0 and h on level 1. Notice that we use $(2t)^{2nb}$ instead of $2tb$ as in [1].

Next, let ρJ_3 be the set of 3-dimensional simplices in $(0, 1] \times \mathbb{C}$ as in Todd [7] where ρ is denoting the grid size. Then ρJ_3 is a refining triangulation of $(0, 1] \times \mathbb{C}$ such that for each $d = 0, 1, 2, \dots$, the set $[2^{-(d+1)}, 2^{-d}] \times \mathbb{C}$ is triangulated and the set $\{2^{-d}\} \times \mathbb{C}$ is subdivided into triangles with diameter all equal to $2^{-d}\sqrt{2}\rho$. The set of facets of all simplices in ρJ_3 is denoted by $(\rho J_3)^2$. For $\sigma \in \rho J_3$ or $\sigma \in (\rho J_3)^2$, the number $t(\sigma)$ denotes the depth of σ in $(0, 1] \times \mathbb{C}$, i.e., $t(\sigma) = \sup\{t | (t, z) \in \sigma\}$. Let $\Phi : [0, 1] \times \mathbb{C} \rightarrow \mathbb{C}$ be the piecewise linear approximation of H with respect to the triangulation ρJ_3 , i.e., let $(t, z) = \sum_{i=1}^4 \lambda_i(t_i, z^i)$ be a point in the simplex in ρJ_3 with vertices (t_i, z^i) for $i = 1, \dots, 4$, so that $\sum_{i=1}^4 \lambda_i = 1$ and $\lambda_i \geq 0$ for $i = 1, \dots, 4$, then

$$\Phi(t, z) = \sum_{i=1}^4 \lambda_i H(t_i, z^i).$$

Let $J(\varepsilon) = \{(s, s^2, \dots, s^n)^T \in \mathbb{R}^n | 0 < s < \varepsilon\}$ for any positive number ε . A 2-dimensional simplex τ in $(\rho J_\varepsilon)^2$ with vertices y^1, y^2, y^3 is complete if $J(\varepsilon)$ is a subset of the convex hull of $H(y^1), H(y^2)$, and $H(y^3)$ for some positive ε . A simplex in ρJ_3 is complete if one of its facets is a complete 2-dimensional simplex. Any complete simplex in ρJ_3 has exactly one pair of complete facets.

The following two propositions are straightforward.

Proposition 2.1 Suppose $\tau \in (\rho J_3)^2$ is complete, then

$$\bar{\tau} \cap \Phi^{-1}(0) \neq \emptyset,$$

where $\bar{\tau}$ is the closure of τ and $\Phi^{-1}(0) = \{(t, z) \in (0, 1] \times \mathbb{C} | \Phi(t, z) = 0\}$ is the zero set of Φ .

Proposition 2.2 Suppose $\tau \in (\rho J_3)^2$ is complete and none of its vertices is a zero of Φ . Then there is at least a pair of its vertices such that their Φ -images span with respect to the origin an angle not less than $2\pi/3$.

The next propositions can be found in [1].

Proposition 2.3 *Suppose that $0 < \rho \leq r_0/(20n)$. Then there are exactly n complete facets in $\{1\} \times \mathbb{C}$.*

To simplify the discussion, in the sequel we consider only the case when $|a_k| \leq 1, k = 1, \dots, n$. Notice that this is only a technical restriction. Moreover, let $|b| \leq 1, \rho = 1$, and $r_0 = 20n$. Then the condition of Proposition 2.3 is satisfied. The next proposition says that all complete simplices lie in a bounded set.

Proposition 2.4 *All complete facets in $(\rho J_3)^2$ lie entirely within a vertical circular cylinder with axis $(0, 1] \times \{0\}$ and radius $R = 40n$.*

The propositions above guarantee that there exist n sequences of adjacent simplices in ρJ_3 having common complete facets. Each sequence starts with some complete facet in $\{1\} \times \mathbb{C}$ and consists of an infinite number of simplices. Since all complete facets lie in a bounded set and the number of complete facets in the set $[2^{-d}, 1] \times \mathbb{C}$ for any integer d is finite, each sequence therefore will reach any level $\{2^{-d}\} \times \mathbb{C}, d = 0, 1, \dots$, within a finite number of simplices. Each sequence of adjacent complete simplices can be followed by complementarity pivoting. Moreover, the sequences lead to the roots of f on level 0, see [1].

Proposition 2.5 *Complementary pivoting procedures which start from the n complete facets in $\{1\} \times \mathbb{C}$ give all n zeros of the polynomial under consideration with the correct multiplicities.*

3 Error estimate

In this section we discuss how far a complete facet in $(\rho J^3)^2$ lies from one of the roots of f when it lies close to the level 0. The next proposition shows how close the roots of two arbitrary polynomials of degree n , lie at most from each other, see [4] for a proof.

Proposition 3.1 *Suppose $f(z) = z^n + \sum_{k=1}^n a_k z^{n-k}$ and $g(z) = z^n + \sum_{k=1}^n b_k z^{n-k}$ are two polynomials of degree n and let $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_n be their zeros, respectively. Let $\mu = \max\{|\alpha_1|, \dots, |\alpha_n|, |\beta_1|, \dots, |\beta_n|\}$ and $\theta = (\sum_{k=1}^n |b_k - a_k| \mu^{n-k})^{1/n}$. Then after appropriate arrangement if necessary, we have that*

$$|\alpha_k - \beta_k| < 2n\theta, \quad k = 1, 2, \dots, n.$$

For any given $t \in (0, \frac{1}{2}]$, let ξ_1, \dots, ξ_k be the zeros of $H(t, z)$, then Proposition 3.1 yields that after appropriate arrangement

$$|\alpha_k - \xi_k| < w(t), k = 1, 2, \dots, n,$$

where

$$w(t) = 2n(2t)^2|b|^{1/n} \leq 2n(2t)^2.$$

Let N be equal to $\lceil \log_2 n \rceil + 2$ where $\lceil \tau \rceil$ stands for the smallest integer not less than τ for any real number τ , then for $t \leq 2^{-N}$ we obtain that $2n2t = 4nt \leq 1$, and thus

$$w(t) \leq 2t.$$

The next proposition can be found in [3].

Proposition 3.2 *For any $w \in \mathbb{C}$ with $|w| < 1$ it holds that*

$$|\arg(1 + w)| \leq \frac{1}{2}\pi|w|.$$

Finally, for any $\xi \in \mathbb{C}$ and $r > 0$, let $B(\xi, r)$ denote the open ball in \mathbb{C} with centre ξ and radius r , i.e., $B(\xi, r) = \{z \in \mathbb{C} | |z - \xi| < r\}$, then the next theorem says that any complete facet τ with depth t at most equal to 2^{-N} lies at most a distance of $(5n + 4)t$ from one of the zeros of f .

Theorem 3.1 . *Let τ be a complete facet in $(\rho J^3)^2$ with vertices y^1, y^2, y^3 and let its depth $t(\tau)$ be less than or equal to 2^{-N} . Then*

$$\tau \subset (0, t(\tau)] \times \bigcup_{k=1}^n B(\alpha_k, (5n + 4)t(\tau)).$$

Proof. We need only to prove the theorem in the case of $n \geq 2$ and $H(y^1), H(y^2)$ and $H(y^3)$ are all unequal to zero. For $i = 1, 2, 3$, let $y^i = (t_i, z^i)$ and let ξ_1^i, \dots, ξ_n^i be the zeros of $H(t_i, \cdot)$ such that these zeros are consistent with the zeros $\alpha_1, \dots, \alpha_n$ of f in the sense of Proposition 3.1. Thus $|\xi_k^i - \alpha_k| \leq 2t_i, i = 1, 2, 3, k = 1, \dots, n$.

In case $t_1 = t_2 = t_3$, we need only to show that the distance of τ and the set of zeros of $H(t_1, \cdot)$ is not greater than $5nt_1$, implying that $\tau \subset \{t_1\} \times B(\alpha_k, (5n + 4)t(\tau))$ for some

$k \in \{1, \dots, n\}$. Suppose that the distance of τ and every ξ_k^1 is greater than $5nt_1$, thus together with Proposition 3.2, for all $i \neq j \in \{1, 2, 3\}$

$$\begin{aligned} \left| \arg \frac{H(t_1, z_i)}{H(t_1, z_j)} \right| &\leq \sum_{k=1}^n \left| \arg \frac{z_i - \xi_k^1}{z_j - \xi_k^1} \right| = \sum_{k=1}^n \left| \arg \left(1 + \frac{z_i - z_j}{z_j - \xi_k^1} \right) \right| \\ &\leq n \frac{\pi}{2} \frac{\sqrt{2}}{5n} < 2\pi/3, \end{aligned}$$

since $|(z_i - z_j)/(z_j - \xi_k^1)| < \sqrt{2} t_1 / (5nt_1) < 1$. This contradicts to the result of Proposition 2.2.

In case not $t_1 = t_2 = t_3$, suppose without loss of generality that $t_1 < t_2$ while $t_1 \leq t_3 \leq t_2$. Now $t(\tau)$ is equal to t_2 . Since the diameter of the simplex τ projected on $\{t_2\} \times \mathbb{C}$ is at most equal to $\sqrt{2} t_2$ we need only to show that the distance of τ and the set of zeros of $H(t_2, \cdot)$ is not greater than $(5n - \sqrt{2})t_2$. Suppose not, then

$$\left| \frac{(z_1 - z_2 + \xi_k^2 - \xi_k^1)}{(z_2 - \xi_k^2)} \right| \leq \frac{\sqrt{2} + 4}{5n - \sqrt{2}} < 1$$

and so

$$\begin{aligned} \left| \arg \frac{H(t_1, z_1)}{H(t_2, z_2)} \right| &\leq \sum_{k=1}^n \left| \arg \frac{z_1 - \xi_k^1}{z_2 - \xi_k^2} \right| = \sum_{k=1}^n \left| \arg \left(1 + \frac{z_1 - z_2 + \xi_k^2 - \xi_k^1}{z_2 - \xi_k^2} \right) \right| \\ &\leq n \frac{\pi}{2} \frac{\sqrt{2} + 4}{5n - \sqrt{2}} < 2\pi/3. \end{aligned}$$

The argument of $|\arg(H(t_1, z_1)/H(t_3, z_3))| < 2\pi/3$ is exactly the same if $t_3 = t_2$ and the same as above if $t_3 = t_1$. Similarly, $|\arg(H(t_3, z_3)/H(t_2, z_2))| < 2\pi/3$ holds. This again contradicts the result of Proposition 2.2. This completes the proof. \square

4 Computational complexity

In this section we discuss how many simplices in the triangulation ρJ_3 will be at most generated before some given accuracy in approximation is reached for computing all zeros of the polynomial f . The calculation of the number of simplices is similar to the one in case of integer labelling.

Proposition 4.1 *Let d be a nonnegative integer and let K be a unit cube in $(0, 1] \times \mathbb{C}$. Let $\xi \in \mathbb{C}$ and Z a disk in \mathbb{C} with ξ as center. Denote s_d as the number of d -dimensional*

simplices in ρJ_3 which meet with $[2^{-(d+1)}, 2^{-d}] \times (K \cap Z)$. Then

$$s_d \leq 14 \text{Vol}(K \cap Z) 2^{2d},$$

where $\text{Vol}(K \cap Z)$ means the volume of $K \cap Z$ in Euclidean two-space.

Proof. This proposition is just a variant of Proposition 3.1 in [3]. \square

The next proposition is a corollary of Theorem 3.1.

Proposition 4.2 *To locate all zeros of f within an accuracy of ε , no computation below level $t = 2^{-D}$ is necessary if ε is small enough, where D is equal to $\log_2[(5n+4)/\varepsilon]$.*

All the propositions above together with Theorem 3.1 imply the following result.

Theorem 4.1 *For $\varepsilon > 0$ small enough, the number of pivots to compute all zeros of f within an accuracy of ε is at most equal to*

$$14\pi[22000n^4 + n(5n+4)^2(D-N)],$$

where $N = \lceil \log_2 n \rceil + 2$ and $D = \lceil \log_2((5n+4)/\varepsilon) \rceil$.

Proof. By Theorem 3.1 and Proposition 2.4, all complete facets lie in either a cylinder of radius $40n$ between the levels $t = 1$ and $t = 2^{-N}$ with center the origin or in n stepped cylinders centred at the zeros of f when $0 < t \leq 2^{-N}$. With Proposition 4.1 the number of complete simplices between $t = 1$ and $t = 2^{-N}$ is at most equal to

$$\begin{aligned} \sum_{i=0}^N 14\pi(40n)^2 2^{2i} &= 14\pi(40n)^2(2^{2N+2} - 1)/3 \\ &\leq 14\pi(4n)^2 2^8 n^2 / 3 \leq 14\pi(22000n^4). \end{aligned}$$

For $d \geq N$, between $t = 2^{-d}$ and $t = 2^{-d+1}$ the number of complete simplices in ρJ_3 is at most equal to

$$n14\pi(5n+4)^2 2^{-2d} 2^{2d} = 14\pi n(5n+4)^2.$$

Hence, by Proposition 4.2, the total number of simplices encountered in computation up to an accuracy of ε is at most equal to

$$14\pi[22000n^4 + n(5n+4)^2(D-N)].$$

The theorem shows that the cost of locating all zeros of f with correct multiplicities and with accuracy equal to ε grows not faster than $O(\max\{n^4, n^3 \log_2(n/\varepsilon)\})$.

References

- [1] Kojima, M., H. Nishino and N. Arima (1979). "A PL homotopy for finding all roots of a polynomial", *Math. Prog.* **16**, 37-62.
- [2] Kuhn, H.W. (1977). "Finding roots of polynomials by pivoting", in *Fixed Points: Algorithms and Applications*, ed. by S. Karamaridiam, Academic Press, 11-39.
- [3] Kuhn, H.W., Z. Wang and S. Xu (1984). "On the cost of computing roots of polynomials", *Math. Prog.* **28**, 156-163.
- [4] Ostrowski, A.M. (1960). *Solution of Equations and Systems of Equations*, 2nd edition, Academic Press.
- [5] Renegar, J. (1985). "On the cost of approximating all roots of a complex polynomial", *Math. Prog.* **32**, 319-336.
- [6] Smale, S. (1981). "The fundamental theorem of algebra and complexity theory", *Bull. Amer. Math. Soc.* **4**, 1-36.
- [7] Todd, M.J. (1976), "On triangulations for computing fixed points", *Math. Prog.* **10**, 322-346.
- [8] Wang, Z. (1989). "A geometrical interpretation of the without-exception feasibility of PL homotopy methods", *Acta Math. Appl. Sinica* **5**, 310-317.
- [9] Wang, Z. and S. Xu (1984). "Approximate zeros and computational complexity", *Scientia Sinica (Series A)* **17**, 566-575.

Discussion Paper Series, CentER, Tilburg University, The Netherlands:

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
9013	A. Horsley and A. Wrobel	The Closedness of the Free-Disposal Hull of a Production Set
9014	A. Horsley and A. Wrobel	The Continuity of the Equilibrium Price Density: The Case of Symmetric Joint Costs, and a Solution to the Shifting-Pattern Problem
9015	A. van den Elzen, G. van der Laan and D. Talman	An Adjustment Process for an Exchange Economy with Linear Production Technologies
9016	P. Deschamps	On Fractional Demand Systems and Budget Share Positivity
9017	B.J. Christensen and N.M. Kiefer	The Exact Likelihood Function for an Empirical Job Search Model
9018	M. Verbeek and Th. Nijman	Testing for Selectivity Bias in Panel Data Models
9019	J.R. Magnus and B. Pesaran	Evaluation of Moments of Ratios of Quadratic Forms in Normal Variables and Related Statistics
9020	A. Robson	Status, the Distribution of Wealth, Social and Private Attitudes to Risk
9021	J.R. Magnus and B. Pesaran	Evaluation of Moments of Quadratic Forms in Normal Variables
9022	K. Kamiya and D. Talman	Linear Stationary Point Problems
9023	W. Emons	Good Times, Bad Times, and Vertical Upstream Integration
9024	C. Dang	The D_2 -Triangulation for Simplicial Homotopy Algorithms for Computing Solutions of Nonlinear Equations
9025	K. Kamiya and D. Talman	Variable Dimension Simplicial Algorithm for Balanced Games
9026	P. Skott	Efficiency Wages, Mark-Up Pricing and Effective Demand
9027	C. Dang and D. Talman	The D_1 -Triangulation in Simplicial Variable Dimension Algorithms for Computing Solutions of Nonlinear Equations
9028	J. Bai, A.J. Jakeman and M. McAleer	Discrimination Between Nested Two- and Three- Parameter Distributions: An Application to Models of Air Pollution

No.	Author(s)	Title
9029	Th. van de Klundert	Crowding out and the Wealth of Nations
9030	Th. van de Klundert and R. Gradus	Optimal Government Debt under Distortionary Taxation
9031	A. Weber	The Credibility of Monetary Target Announce- ments: An Empirical Evaluation
9032	J. Osiewalski and M. Steel	Robust Bayesian Inference in Elliptical Regression Models
9033	C. R. Wichers	The Linear-Algebraic Structure of Least Squares
9034	C. de Vries	On the Relation between GARCH and Stable Processes
9035	M.R. Baye, D.W. Jansen and Q. Li	Aggregation and the "Random Objective" Justification for Disturbances in Complete Demand Systems
9036	J. Driffill	The Term Structure of Interest Rates: Structural Stability and Macroeconomic Policy Changes in the UK
9037	F. van der Ploeg	Budgetary Aspects of Economic and Monetary Integration in Europe
9038	A. Robson	Existence of Nash Equilibrium in Mixed Strategies for Games where Payoffs Need not Be Continuous in Pure Strategies
9039	A. Robson	An "Informationally Robust Equilibrium" for Two-Person Nonzero-Sum Games
9040	M.R. Baye, G. Tian and J. Zhou	The Existence of Pure-Strategy Nash Equilibrium in Games with Payoffs that are not Quasiconcave
9041	M. Burnovsky and I. Zang	"Costless" Indirect Regulation of Monopolies with Substantial Entry Cost
9042	P.J. Deschamps	Joint Tests for Regularity and Autocorrelation in Allocation Systems
9043	S. Chib, J. Osiewalski and M. Steel	Posterior Inference on the Degrees of Freedom Parameter in Multivariate-t Regression Models
9044	H.A. Keuzenkamp	The Probability Approach in Economic Method- ology: On the Relation between Haavelmo's Legacy and the Methodology of Economics
9045	I.M. Bomze and E.E.C. van Damme	A Dynamical Characterization of Evolution- arily Stable States
9046	E. van Damme	On Dominance Solvable Games and Equilibrium Selection Theories

No.	Author(s)	Title
9047	J. Driffill	Changes in Regime and the Term Structure: A Note
9048	A.J.J. Talman	General Equilibrium Programming
9049	H.A. Keuzenkamp and F. van der Ploeg	Saving, Investment, Government Finance and the Current Account: The Dutch Experience
9050	C. Dang and A.J.J. Talman	The D _n -Triangulation in Simplicial Variable Dimension Algorithms on the Unit Simplex for Computing Fixed Points
9051	M. Baye, D. Kovenock and C. de Vries	The All-Pay Auction with Complete Information
9052	H. Carlsson and E. van Damme	Global Games and Equilibrium Selection
9053	M. Baye and D. Kovenock	How to Sell a Pickup Truck: "Beat-or-Pay" Advertisements as Facilitating Devices
9054	Th. van de Klundert	The Ultimate Consequences of the New Growth Theory; An Introduction to the Views of M. Fitzgerald Scott
9055	P. Kooreman	Nonparametric Bounds on the Regression Coefficients when an Explanatory Variable is Categorized
9056	R. Bartels and D.G. Fiebig	Integrating Direct Metering and Conditional Demand Analysis for Estimating End-Use Loads
9057	M.R. Veall and K.F. Zimmermann	Evaluating Pseudo-R ² 's for Binary Probit Models
9058	R. Bartels and D.G. Fiebig	More on the Grouped Heteroskedasticity Model
9059	F. van der Ploeg	Channels of International Policy Transmission
9060	H. Bester	The Role of Collateral in a Model of Debt Renegotiation
9061	F. van der Ploeg	Macroeconomic Policy Coordination during the Various Phases of Economic and Monetary Integration in Europe
9062	E. Bennett and E. van Damme	Demand Commitment Bargaining: - The Case of Apex Games
9063	S. Chib, J. Osiewalski and M. Steel	Regression Models under Competing Covariance Matrices: A Bayesian Perspective
9064	M. Verbeek and Th. Nijman	Can Cohort Data Be Treated as Genuine Panel Data?

No.	Author(s)	Title
9065	F. van der Ploeg and A. de Zeeuw	International Aspects of Pollution Control
9066	F.C. Drost and Th. E. Nijman	Temporal Aggregation of GARCH Processes
9067	Y. Dai and D. Talman	Linear Stationary Point Problems on Unbounded Polyhedra
9068	Th. Nijman and R. Beetsma	Empirical Tests of a Simple Pricing Model for Sugar Futures
9069	F. van der Ploeg	Short-Sighted Politicians and Erosion of Government Assets
9070	E. van Damme	Fair Division under Asymmetric Information
9071	J. Eichberger, H. Haller and F. Milne	Naive Bayesian Learning in 2 x 2 Matrix Games
9072	G. Alogoskoufis and F. van der Ploeg	Endogenous Growth and Overlapping Generations
9073	K.C. Fung	Strategic Industrial Policy for Cournot and Bertrand Oligopoly: Management-Labor Cooperation as a Possible Solution to the Market Structure Dilemma
9101	A. van Soest	Minimum Wages, Earnings and Employment
9102	A. Barten and M. McAleer	Comparing the Empirical Performance of Alternative Demand Systems
9103	A. Weber	EMS Credibility
9104	G. Alogoskoufis and F. van der Ploeg	Debts, Deficits and Growth in Interdependent Economies
9105	R.M.W.J. Beetsma	Bands and Statistical Properties of EMS Exchange Rates
9106	C.N. Teulings	The Diverging Effects of the Business Cycle on the Expected Duration of Job Search
9107	E. van Damme	Refinements of Nash Equilibrium
9108	E. van Damme	Equilibrium Selection in 2 x 2 Games
9109	G. Alogoskoufis and F. van der Ploeg	Money and Growth Revisited
9110	L. Samuelson	Dominated Strategies and Common Knowledge
9111	F. van der Ploeg and Th. van de Klundert	Political Trade-off between Growth and Government Consumption

No.	Author(s)	Title
9112	Th. Nijman, F. Palm and C. Wolff	Premia in Forward Foreign Exchange as Unobserved Components
9113	H. Bester	Bargaining vs. Price Competition in a Market with Quality Uncertainty
9114	R.P. Gilles, G. Owen and R. van den Brink	Games with Permission Structures: The Conjunctive Approach
9115	F. van der Ploeg	Unanticipated Inflation and Government Finance: The Case for an Independent Common Central Bank
9116	N. Rankin	Exchange Rate Risk and Imperfect Capital Mobility in an Optimising Model
9117	E. Bomhoff	Currency Convertibility: When and How? A Contribution to the Bulgarian Debate!
9118	E. Bomhoff	Stability of Velocity in the G-7 Countries: A Kalman Filter Approach
9119	J. Osiewalski and M. Steel	Bayesian Marginal Equivalence of Elliptical Regression Models
9120	S. Bhattacharya, J. Glazer and D. Sappington	Licensing and the Sharing of Knowledge in Research Joint Ventures
9121	J.W. Friedman and L. Samuelson	An Extension of the "Folk Theorem" with Continuous Reaction Functions
9122	S. Chib, J. Osiewalski and M. Steel	A Bayesian Note on Competing Correlation Structures in the Dynamic Linear Regression Model
9123	Th. van de Klundert and L. Meijdam	Endogenous Growth and Income Distribution
9124	S. Bhattacharya	Banking Theory: The Main Ideas
9125	J. Thomas	Non-Computable Rational Expectations Equilibria
9126	J. Thomas and T. Worrall	Foreign Direct Investment and the Risk of Expropriation
9127	T. Gao, A.J.J. Talman and Z. Wang	Modification of the Kojima-Nishino-Arima Algorithm and its Computational Complexity